

INVARIANTS OF LINES ON SURFACES IN $\mathbb{R}^4$

GEORGI GANCHEV AND VELICHKA MILOUSHEVA

ABSTRACT. Considering the tangent plane at a point to a surface in the four-dimensional Euclidean space, we find an invariant of a pair of two tangents in this plane. If this invariant is zero, the two tangents are said to be conjugate. When the two tangents coincide with a given tangent, then we obtain the normal curvature of this tangent. Asymptotic tangents (curves) are characterized by zero normal curvature. Considering the invariant of the pair of a given tangent and its orthogonal one, we introduce the geodesic torsion of this tangent. We obtain that principal tangents (curves) are characterized by zero geodesic torsion.

The invariants $\varkappa$ and k are introduced as the symmetric functions of the two principal normal curvatures. The geometric meaning of the semi-sum $\varkappa$ of the principal normal curvatures is equal (up to a sign) to the curvature of the normal connection of the surface. The number of asymptotic tangents at a point of the surface is determined by the sign of the invariant k . In the case $k = 0$ there exists a one-parameter family of asymptotic lines, which are principal. We find examples of such surfaces ($k = 0$) in the class of the general rotational surfaces (in the sense of Moore). The principal asymptotic lines on these surfaces are helices in the four-dimensional Euclidean space.

1. PRELIMINARIES

In [1] we introduced a linear map γ of Weingarten type in the tangent space at any point of a surface in the four-dimensional space $\mathbb{R}^4$. Analogously to the classical theory of surfaces in $\mathbb{R}^3$ the map γ generates second fundamental form II on the surface.

In the present paper we introduce two invariants of a line on a surface M^2 in $\mathbb{R}^4$ in terms of an invariant ζ_{g_1, g_2} of two tangents g_1, g_2 at a point of M^2 . Here we define *conjugate tangents* by the condition $\zeta_{g_1, g_2} = 0$. We prove that the conjugacy in terms of ζ_{g_1, g_2} is the conjugacy with respect to the second fundamental form. Thus we obtain a geometric interpretation of the second fundamental form and the Weingarten map of the surface.

Let $M^2 : z = z(u, v)$, $(u, v) \in \mathcal{D}$ ($\mathcal{D} \subset \mathbb{R}^2$) be a 2-dimensional surface in $\mathbb{R}^4$ with tangent space $T_p M^2 = \text{span}\{z_u, z_v\}$ at any point $p \in M^2$, and E, F, G be the coefficients of the first fundamental form, $W = \sqrt{EG - F^2}$. We choose an orthonormal normal frame field $\{e_1, e_2\}$ of M^2 so that the quadruple $\{z_u, z_v, e_1, e_2\}$ is positive oriented in $\mathbb{R}^4$. Then the following derivative formulas hold:

$$\begin{aligned}\nabla'_{z_u} z_u &= z_{uu} = \Gamma_{11}^1 z_u + \Gamma_{11}^2 z_v + c_{11}^1 e_1 + c_{11}^2 e_2, \\ \nabla'_{z_u} z_v &= z_{uv} = \Gamma_{12}^1 z_u + \Gamma_{12}^2 z_v + c_{12}^1 e_1 + c_{12}^2 e_2, \\ \nabla'_{z_v} z_v &= z_{vv} = \Gamma_{22}^1 z_u + \Gamma_{22}^2 z_v + c_{22}^1 e_1 + c_{22}^2 e_2,\end{aligned}$$

where Γ_{ij}^k are the Christoffel's symbols and c_{ij}^k , $i, j, k = 1, 2$ are functions on M^2 .

Denoting by σ the second fundamental tensor of M^2 , we have

$$\begin{aligned}(1.1) \quad \sigma(z_u, z_u) &= c_{11}^1 e_1 + c_{11}^2 e_2, \\ \sigma(z_u, z_v) &= c_{12}^1 e_1 + c_{12}^2 e_2, \\ \sigma(z_v, z_v) &= c_{22}^1 e_1 + c_{22}^2 e_2.\end{aligned}$$

2000 *Mathematics Subject Classification*. Primary 53A07, Secondary 53A10.

Key words and phrases. Curves on surfaces in the four-dimensional Euclidean space, normal curvature, asymptotic lines, geodesic torsion, principal lines.

The three pairs of normal vectors $\{\sigma(z_u, z_u), \sigma(z_u, z_v)\}$, $\{\sigma(z_u, z_u), \sigma(z_v, z_v)\}$, $\{\sigma(z_u, z_v), \sigma(z_v, z_v)\}$ form three parallelograms with oriented areas

$$\Delta_1 = \begin{vmatrix} c_{11}^1 & c_{12}^1 \\ c_{11}^2 & c_{12}^2 \end{vmatrix}, \quad \Delta_2 = \begin{vmatrix} c_{11}^1 & c_{22}^1 \\ c_{11}^2 & c_{22}^2 \end{vmatrix}, \quad \Delta_3 = \begin{vmatrix} c_{12}^1 & c_{22}^1 \\ c_{12}^2 & c_{22}^2 \end{vmatrix},$$

respectively. These oriented areas determine three functions $L = \frac{2\Delta_1}{W}$, $M = \frac{\Delta_2}{W}$, $N = \frac{2\Delta_3}{W}$, which change in the same way as the coefficients E, F, G under any change of the parameters (u, v) .

Similarly to the theory of surfaces in $\mathbb{R}^3$, using the functions E, F, G and L, M, N in [1] we introduced the linear map γ in the tangent space at any point of M^2

$$\gamma : T_p M^2 \rightarrow T_p M^2,$$

defined by the equalities

$$\begin{aligned} \gamma(z_u) &= \gamma_1^1 z_u + \gamma_1^2 z_v, \\ \gamma(z_v) &= \gamma_2^1 z_u + \gamma_2^2 z_v, \end{aligned}$$

where

$$\gamma_1^1 = \frac{FM - GL}{EG - F^2}, \quad \gamma_1^2 = \frac{FL - EM}{EG - F^2}, \quad \gamma_2^1 = \frac{FN - GM}{EG - F^2}, \quad \gamma_2^2 = \frac{FM - EN}{EG - F^2}.$$

The linear map γ of Weingarten type at the point $p \in M^2$ is invariant with respect to changes of parameters on M^2 as well as to motions in $\mathbb{R}^4$. This implies that the functions

$$k = \frac{LN - M^2}{EG - F^2}, \quad \varkappa = \frac{EN + GL - 2FM}{2(EG - F^2)}$$

are invariants of the surface M^2 .

The invariant $\varkappa$ turns out to be the curvature of the normal connection of the surface M^2 in $\mathbb{R}^4$.

As in the classical case, the invariants k and $\varkappa$ divide the points of M^2 into four types: flat, elliptic, parabolic and hyperbolic. The surfaces consisting of flat points satisfy the conditions

$$k(u, v) = 0, \quad \varkappa(u, v) = 0, \quad (u, v) \in \mathcal{D},$$

or equivalently $L(u, v) = 0$, $M(u, v) = 0$, $N(u, v) = 0$, $(u, v) \in \mathcal{D}$. These surfaces are either planar surfaces (there exists a hyperplane $\mathbb{R}^3 \subset \mathbb{R}^4$ containing M^2) or developable ruled surfaces in $\mathbb{R}^4$ [1].

Further we consider surfaces free of flat points, i.e. $(L, M, N) \neq (0, 0, 0)$.

Let $X = \lambda z_u + \mu z_v$, $(\lambda, \mu) \neq (0, 0)$ be a tangent vector at a point $p \in M^2$. The map γ determines a second fundamental form of the surface M^2 at p as follows:

$$II(\lambda, \mu) = -g(\gamma(X), X) = L\lambda^2 + 2M\lambda\mu + N\mu^2, \quad \lambda, \mu \in \mathbb{R}.$$

As in the classical differential geometry of surfaces in $\mathbb{R}^3$ the second fundamental form II determines conjugate tangents at a point p of M^2 . Two tangents $g_1 : X_1 = \lambda_1 z_u + \mu_1 z_v$ and $g_2 : X_2 = \lambda_2 z_u + \mu_2 z_v$ are said to be *conjugate tangents* if $II(\lambda_1, \mu_1; \lambda_2, \mu_2) = 0$, i.e.

$$L\lambda_1\lambda_2 + M(\lambda_1\mu_2 + \lambda_2\mu_1) + N\mu_1\mu_2 = 0.$$

In the next section we shall introduce conjugate tangents in a geometric way.

2. INVARIANTS OF A TANGENT ON A SURFACE IN $\mathbb{R}^4$

Let g be a tangent at the point $p \in M^2$ determined by the vector $X = \lambda z_u + \mu z_v$. We consider the map $\sigma_g : T_p M^2 \rightarrow (T_p M^2)^\perp$, defined by

$$(2.1) \quad \sigma_g(Y) = \sigma \left(\frac{\lambda z_u + \mu z_v}{\sqrt{I(\lambda, \mu)}}, Y \right), \quad Y \in T_p M^2.$$

Obviously σ_g is a linear map, which does not depend on the choice of the vector X collinear with g . Using (1.1) and (2.1) we obtain the following decomposition of the normal vectors $\sigma_g(z_u)$ and $\sigma_g(z_v)$:

$$(2.2) \quad \begin{aligned} \sigma_g(z_u) &= \frac{\lambda c_{11}^1 + \mu c_{12}^1}{\sqrt{I(\lambda, \mu)}} e_1 + \frac{\lambda c_{11}^2 + \mu c_{12}^2}{\sqrt{I(\lambda, \mu)}} e_2, \\ \sigma_g(z_v) &= \frac{\lambda c_{12}^1 + \mu c_{22}^1}{\sqrt{I(\lambda, \mu)}} e_1 + \frac{\lambda c_{12}^2 + \mu c_{22}^2}{\sqrt{I(\lambda, \mu)}} e_2. \end{aligned}$$

Let $g_1 : X_1 = \lambda_1 z_u + \mu_1 z_v$ and $g_2 : X_2 = \lambda_2 z_u + \mu_2 z_v$ be two tangents at the point $p \in M^2$. The oriented areas of the parallelograms determined by the pairs of normal vectors $\sigma_{g_1}(z_u), \sigma_{g_2}(z_v)$ and $\sigma_{g_2}(z_u), \sigma_{g_1}(z_v)$ are denoted by $S(\sigma_{g_1}(z_u), \sigma_{g_2}(z_v))$, and $S(\sigma_{g_2}(z_u), \sigma_{g_1}(z_v))$, respectively. We assign the quantity ζ_{g_1, g_2} to the pair of tangents g_1, g_2 , defined by

$$(2.3) \quad \zeta_{g_1, g_2} = \frac{S(\sigma_{g_1}(z_u), \sigma_{g_2}(z_v))}{W} + \frac{S(\sigma_{g_2}(z_u), \sigma_{g_1}(z_v))}{W}.$$

Proposition 2.1. *The quantity ζ_{g_1, g_2} is invariant under any change of the parameters on M^2 .*

Proof: Using equalities (2.2) we calculate that

$$\zeta_{g_1, g_2} = \frac{L\lambda_1\lambda_2 + M(\lambda_1\mu_2 + \mu_1\lambda_2) + N\mu_1\mu_2}{\sqrt{I(\lambda_1, \mu_1)}\sqrt{I(\lambda_2, \mu_2)}} = \frac{II(\lambda_1, \mu_1; \lambda_2, \mu_2)}{\sqrt{I(\lambda_1, \mu_1)}\sqrt{I(\lambda_2, \mu_2)}}.$$

Now, let

$$\begin{aligned} u &= u(\bar{u}, \bar{v}); \\ v &= v(\bar{u}, \bar{v}), \quad (\bar{u}, \bar{v}) \in \bar{\mathcal{D}}, \quad \bar{\mathcal{D}} \subset \mathbb{R}^2 \end{aligned}$$

be a smooth change of the parameters (u, v) on M^2 with $J = u_{\bar{u}} v_{\bar{v}} - u_{\bar{v}} v_{\bar{u}} \neq 0$. Then

$$\begin{aligned} z_{\bar{u}} &= z_u u_{\bar{u}} + z_v v_{\bar{u}}, \\ z_{\bar{v}} &= z_u u_{\bar{v}} + z_v v_{\bar{v}} \end{aligned}$$

and $\bar{E}\bar{G} - \bar{F}^2 = J^2(EG - F^2)$, hence $\bar{W} = \varepsilon J W$, $\varepsilon = \text{sign } J$. The functions E, F, G and L, M, N change as follows under the change of the parametrization:

$$\begin{aligned} \bar{E} &= u_{\bar{u}}^2 E + 2 u_{\bar{u}} v_{\bar{u}} F + v_{\bar{u}}^2 G, & \bar{L} &= \varepsilon(u_{\bar{u}}^2 L + 2 u_{\bar{u}} v_{\bar{u}} M + v_{\bar{u}}^2 N), \\ \bar{F} &= u_{\bar{u}} u_{\bar{v}} E + (u_{\bar{u}} v_{\bar{v}} + v_{\bar{u}} u_{\bar{v}}) F + v_{\bar{u}} v_{\bar{v}} G, & \bar{M} &= \varepsilon(u_{\bar{u}} u_{\bar{v}} L + (u_{\bar{u}} v_{\bar{v}} + v_{\bar{u}} u_{\bar{v}}) M + v_{\bar{u}} v_{\bar{v}} N) \\ \bar{G} &= u_{\bar{v}}^2 E + 2 u_{\bar{v}} v_{\bar{v}} F + v_{\bar{v}}^2 G, & \bar{N} &= \varepsilon(u_{\bar{v}}^2 L + 2 u_{\bar{v}} v_{\bar{v}} M + v_{\bar{v}}^2 N). \end{aligned}$$

If $X = \lambda z_u + \mu z_v = \bar{\lambda} z_{\bar{u}} + \bar{\mu} z_{\bar{v}}$, then $\lambda = u_{\bar{u}} \bar{\lambda} + u_{\bar{v}} \bar{\mu}$, $\mu = v_{\bar{u}} \bar{\lambda} + v_{\bar{v}} \bar{\mu}$.

The first and the second fundamental forms change as follows:

$$\bar{I}(\bar{\lambda}, \bar{\mu}) = I(\lambda, \mu); \quad \bar{II}(\bar{\lambda}, \bar{\mu}) = \varepsilon II(\lambda, \mu).$$

Hence,

$$\bar{\zeta}_{g_1, g_2} = \frac{\bar{II}(\bar{\lambda}_1, \bar{\mu}_1; \bar{\lambda}_2, \bar{\mu}_2)}{\sqrt{\bar{I}(\bar{\lambda}_1, \bar{\mu}_1)}\sqrt{\bar{I}(\bar{\lambda}_2, \bar{\mu}_2)}} = \frac{\varepsilon II(\lambda_1, \mu_1; \lambda_2, \mu_2)}{\sqrt{I(\lambda_1, \mu_1)}\sqrt{I(\lambda_2, \mu_2)}} = \varepsilon \zeta_{g_1, g_2}.$$

Consequently, ζ_{g_1, g_2} is invariant (up to the orientation of the tangent space or the normal space). $\square$

Definition 2.2. Two tangents $g_1 : X_1 = \lambda_1 z_u + \mu_1 z_v$ and $g_2 : X_2 = \lambda_2 z_u + \mu_2 z_v$ are said to be *conjugate tangents* if $\zeta_{g_1, g_2} = 0$.

Obviously, $\zeta_{g_1, g_2} = 0$ if and only if $L\lambda_1\lambda_2 + M(\lambda_1\mu_2 + \lambda_2\mu_1) + N\mu_1\mu_2 = 0$. Hence, the tangents g_1 and g_2 are conjugate according to Definition 2.2 if and only if they are conjugate with respect to the second fundamental form II .

We shall assign two invariants ν_g and α_g to any tangent g of the surface in the following way. Let $g : X = \lambda z_u + \mu z_v$ be a tangent and $g^\perp$ be its orthogonal tangent, determined by the vector

$$(2.4) \quad X^\perp = -\frac{F\lambda + G\mu}{W} z_u + \frac{E\lambda + F\mu}{W} z_v.$$

We define

$$\nu_g = \zeta_{g, g^\perp}; \quad \alpha_g = \zeta_{g, g^\perp}.$$

We call ν_g the *normal curvature* of the tangent g , and α_g - the *geodesic torsion* of g .

The equality (2.3) implies that

$$\nu_g = 2 \frac{S(\sigma_g(z_u), \sigma_g(z_v))}{W} = \frac{II(\lambda, \mu)}{I(\lambda, \mu)}.$$

Hence, the normal curvature of the tangent g is two times the oriented area of the parallelogram determined by the normal vectors $\sigma_g(z_u)$ and $\sigma_g(z_v)$. The invariant ν_g is expressed by the first and the second fundamental forms of the surface in the same way as the normal curvature of a tangent in the theory of surfaces in $\mathbb{R}^3$.

Using (2.3) and (2.4) we get

$$\alpha_g = \frac{\lambda^2(EM - FL) + \lambda\mu(EN - GL) + \mu^2(FN - GM)}{WI(\lambda, \mu)}.$$

The last formula shows that α_g is expressed by the coefficients of the first and the second fundamental forms in the same way as the geodesic torsion in the theory of surfaces in $\mathbb{R}^3$.

Definition 2.3. A tangent $g : X = \lambda z_u + \mu z_v$ is said to be *asymptotic* if it is self-conjugate.

Proposition 2.4. A tangent g is asymptotic if and only if $\nu_g = 0$.

Definition 2.5. A tangent $g : X = \lambda z_u + \mu z_v$ is said to be *principal* if it is perpendicular to its conjugate.

Proposition 2.6. A tangent g is principal if and only if $\alpha_g = 0$.

The equation for the asymptotic tangents at a point $p \in M^2$ is

$$L\lambda^2 + 2M\lambda\mu + N\mu^2 = 0.$$

If p is an elliptic point of M^2 ($k > 0$) then there are no asymptotic tangents through p ; if p is a hyperbolic point ($k < 0$) then there are two asymptotic tangents passing through p , and if p is a parabolic point ($k = 0$) then there is one asymptotic tangent through p .

A line $c : u = u(q), v = v(q); q \in J \subset \mathbb{R}$ on M^2 is said to be an *asymptotic line* if its tangent at any point is asymptotic.

The equation for the principal tangents at a point $p \in M^2$ is

$$\begin{vmatrix} E & F \\ L & M \end{vmatrix} \lambda^2 + \begin{vmatrix} E & G \\ L & N \end{vmatrix} \lambda\mu + \begin{vmatrix} F & G \\ M & N \end{vmatrix} \mu^2 = 0.$$

A line $c : u = u(q), v = v(q); q \in J \subset \mathbb{R}$ on M^2 is said to be a *principal line* (a *line of curvature*) if its tangent at any point is principal. The surface M^2 is parameterized by the principal lines if and only if $F = 0, M = 0$.

The normal curvatures $\nu' = \frac{L}{E}$ and $\nu'' = \frac{N}{G}$ of the principal tangents are said to be *principal normal curvatures* of M^2 . The invariants k and κ of M^2 are expressed by the principal normal curvatures ν' and ν'' as follows:

$$k = \nu'\nu''; \quad \kappa = \frac{\nu' + \nu''}{2}.$$

3. EXAMPLES OF SURFACES WITH $k = 0$

In this section we consider general (in the sense of C. Moore) rotational surfaces in $\mathbb{R}^4$ whose meridians lie in two-dimensional planes. We shall find all such surfaces consisting of parabolic points.

Considering general rotations in $\mathbb{R}^4$, C. Moore introduced general rotational surfaces [2] (see also [3, 4]). In the case when the meridians lie in two-dimensional planes the general rotational surface can be parameterized as follows:

$$M^2 : z(u, v) = (f(u) \cos \alpha v, f(u) \sin \alpha v, g(u) \cos \beta v, g(u) \sin \beta v); \quad u \in J \subset \mathbb{R}, v \in [0; 2\pi),$$

where $f(u)$ and $g(u)$ are smooth functions, satisfying $\alpha^2 f^2(u) + \beta^2 g^2(u) > 0$, $f'^2(u) + g'^2(u) > 0$, $u \in J$, and α, β are positive constants.

Each parametric curve $u = u_0 = \text{const}$ of M^2 is given by

$$c_v : z(v) = (a \cos \alpha v, a \sin \alpha v, b \cos \beta v, b \sin \beta v); \quad a = f(u_0), b = g(u_0)$$

and its Frenet curvatures are $\kappa_{c_v} = \sqrt{\frac{a^2 \alpha^4 + b^2 \beta^4}{a^2 \alpha^2 + b^2 \beta^2}}$; $\tau_{c_v} = \frac{ab\alpha\beta(\alpha^2 - \beta^2)}{\sqrt{a^2 \alpha^4 + b^2 \beta^4} \sqrt{a^2 \alpha^2 + b^2 \beta^2}}$; $\sigma_{c_v} = \frac{\alpha\beta \sqrt{a^2 \alpha^2 + b^2 \beta^2}}{\sqrt{a^2 \alpha^4 + b^2 \beta^4}}$. Hence, when $\alpha \neq \beta$ each parametric curve $u = \text{const}$ is a curve in $\mathbb{R}^4$ with constant curvatures (helix), and when $\alpha = \beta$ each parametric curve $u = \text{const}$ is a circle.

Each parametric curve $v = v_0 = \text{const}$ of M^2 is given by

$$c_u : z(u) = (A_1 f(u), A_2 f(u), B_1 g(u), B_2 g(u)),$$

where $A_1 = \cos \alpha v_0$, $A_2 = \sin \alpha v_0$, $B_1 = \cos \beta v_0$, $B_2 = \sin \beta v_0$. The Frenet curvatures of c_u are $\kappa_{c_u} = \frac{|g'f'' - f'g''|}{(\sqrt{f'^2 + g'^2})^3}$; $\tau_{c_u} = 0$. Hence, c_u is a plane curve with curvature $\kappa_{c_u} = \frac{|g'f'' - f'g''|}{(\sqrt{f'^2 + g'^2})^3}$. So, for each $v = \text{const}$ the parametric curves c_u are congruent in $\mathbb{R}^4$. These curves are called *meridians* of M^2 .

The tangent space of M^2 is spanned by the vector fields

$$\begin{aligned} z_u &= (f' \cos \alpha v, f' \sin \alpha v, g' \cos \beta v, g' \sin \beta v); \\ z_v &= (-\alpha f \sin \alpha v, \alpha f \cos \alpha v, -\beta g \sin \beta v, \beta g \cos \beta v). \end{aligned}$$

Hence, the coefficients of the first fundamental form are $E = f'^2(u) + g'^2(u)$; $F = 0$; $G = \alpha^2 f^2(u) + \beta^2 g^2(u)$. We consider the following orthonormal tangent frame field

$$\begin{aligned} x &= \frac{1}{\sqrt{f'^2 + g'^2}} (f' \cos \alpha v, f' \sin \alpha v, g' \cos \beta v, g' \sin \beta v); \\ y &= \frac{1}{\sqrt{\alpha^2 f^2 + \beta^2 g^2}} (-\alpha f \sin \alpha v, \alpha f \cos \alpha v, -\beta g \sin \beta v, \beta g \cos \beta v); \end{aligned}$$

and the following orthonormal normal frame field

$$n_1 = \frac{1}{\sqrt{f'^2 + g'^2}} (g' \cos \alpha v, g' \sin \alpha v, -f' \cos \beta v, -f' \sin \beta v);$$

$$n_2 = \frac{1}{\sqrt{\alpha^2 f^2 + \beta^2 g^2}} (-\beta g \sin \alpha v, \beta g \cos \alpha v, \alpha f \sin \beta v, -\alpha f \cos \beta v).$$

$\{x, y, n_1, n_2\}$ is a positive oriented orthonormal frame field in $\mathbb{R}^4$.

We calculate the functions c_{ij}^k , $i, j, k = 1, 2$ and find the coefficients L , M and N of the second fundamental form of M^2 :

$$L = \frac{2\alpha\beta(gf' - fg')(g'f'' - f'g'')}{(\alpha^2 f^2 + \beta^2 g^2)(f'^2 + g'^2)}; \quad M = 0; \quad N = \frac{-2\alpha\beta(gf' - fg')(\beta^2 gf' - \alpha^2 fg')}{(\alpha^2 f^2 + \beta^2 g^2)(f'^2 + g'^2)}.$$

Consequently, the invariants k , $\varkappa$ and the Gauss curvature K of M^2 are expressed as follows:

$$k = \frac{-4\alpha^2\beta^2(gf' - fg')^2(g'f'' - f'g'')(\beta^2 gf' - \alpha^2 fg')}{(\alpha^2 f^2 + \beta^2 g^2)^3(f'^2 + g'^2)^3};$$

$$\varkappa = \frac{\alpha\beta(gf' - fg')}{(\alpha^2 f^2 + \beta^2 g^2)^2(f'^2 + g'^2)^2} ((\alpha^2 f^2 + \beta^2 g^2)(g'f'' - f'g'') - (f'^2 + g'^2)(\beta^2 gf' - \alpha^2 fg'));$$

$$K = \frac{(\alpha^2 f^2 + \beta^2 g^2)(\beta^2 gf' - \alpha^2 fg')(g'f'' - f'g'') - \alpha^2\beta^2(f'^2 + g'^2)(gf' - fg')^2}{(\alpha^2 f^2 + \beta^2 g^2)^2(f'^2 + g'^2)^2}.$$

Note that the surface M^2 is parameterized by the principal lines ($F = M = 0$).

Now we shall find the general rotational surfaces with $k = 0$. Without loss of generality we assume that the meridian m is defined by $f = u$; $g = g(u)$. Then

$$k = \frac{4\alpha^2\beta^2(g - ug')^2g''(\beta^2 g - \alpha^2 ug')}{(\alpha^2 u^2 + \beta^2 g^2)^3(1 + g'^2)^3}.$$

The invariant k is zero in the following three cases:

1. $g(u) = au$, $a = \text{const} \neq 0$. In this case $k = \varkappa = K = 0$, and M^2 is a developable surface in $\mathbb{R}^4$.

2. $g(u) = au + b$, $a = \text{const} \neq 0, b = \text{const} \neq 0$. In this case $k = 0$, but $\varkappa \neq 0$, $K \neq 0$. Consequently, M^2 is a non-developable ruled surface in $\mathbb{R}^4$.

3. $g(u) = cu^{\frac{\beta^2}{\alpha^2}}$, $c = \text{const} \neq 0$. In case of $\alpha \neq \beta$ we get $k = 0$, and the invariants $\varkappa$ and K are given by

$$\varkappa = \frac{c^2\alpha^{-5}\beta^3(\beta^2 - \alpha^2)^2u^{2\frac{\beta^2 - \alpha^2}{\alpha^2}}}{\left(\alpha^2 u^2 + \beta^2 c^2 u^{2\frac{\beta^2}{\alpha^2}}\right) \left(1 + c^2 \frac{\beta^4}{\alpha^4} u^{2\frac{\beta^2 - \alpha^2}{\alpha^2}}\right)^2}; \quad K = -\frac{c^2\alpha^{-2}\beta^2(\beta^2 - \alpha^2)^2u^{2\frac{\beta^2}{\alpha^2}}}{\left(\alpha^2 u^2 + \beta^2 c^2 u^{2\frac{\beta^2}{\alpha^2}}\right)^2 \left(1 + c^2 \frac{\beta^4}{\alpha^4} u^{2\frac{\beta^2 - \alpha^2}{\alpha^2}}\right)}.$$

Hence, $\varkappa \neq 0$, $K \neq 0$. In this case the parametric lines $u = \text{const}$ and $v = \text{const}$ are not straight lines. This is the most interesting example of general rotational surfaces with $k = 0$.

Since $k = 0$, one of the principal normal curvatures of M^2 is zero ($\nu'' = \frac{N}{G} = 0$). Hence, the parametric v -lines of M^2 are asymptotic principal lines. Moreover, these lines are helices in $\mathbb{R}^4$.

Acknowledgements: The second author is partially supported by "L. Karavelov" Civil Engineering Higher School, Sofia, Bulgaria under Contract No 10/2010.

REFERENCES

- [1] Ganchev G. and V. Milousheva. *On the theory of surfaces in the four-dimensional Euclidean space*. Kodai Math. J., **31** (2008), 183-198.
- [2] Moore C. *Surfaces of rotation in a space of four dimensions*. The Annals of Math., 2nd Ser., **21** (1919), 2, 81-93.
- [3] Moore C. and E. Wilson. *A general theory of surfaces*. J. Nat. Acad. Proc. **2** (1916), 273-278.
- [4] Moore C. and E. Wilson. *Differential geometry of two-dimensional surfaces in hyperspaces*. Proc. Acad. Arts Sci. **52** (1916), 267-368.

BULGARIAN ACADEMY OF SCIENCES, INSTITUTE OF MATHEMATICS AND INFORMATICS, ACAD. G. BONCHEV
STR. BL. 8, 1113 SOFIA, BULGARIA

E-mail address: `ganchev@math.bas.bg`

BULGARIAN ACADEMY OF SCIENCES, INSTITUTE OF MATHEMATICS AND INFORMATICS, ACAD. G. BONCHEV
STR. BL. 8, 1113, SOFIA, BULGARIA; "L. KARAVELOV" CIVIL ENGINEERING HIGHER SCHOOL, 175
SUHODOLSKA STR., 1373 SOFIA, BULGARIA

E-mail address: `vmil@math.bas.bg`